ELSEVIER

International Journal of Pharmaceutics 122 (1995) 131

international journal of pharmaceutics

Author Index

Adler, J., see Aspden, T.J. (122) 69
Al-Shamkhani, A. and Duncan, R.
Synthesis, controlled release properties and antitumour activity of alginate-*cis*-aconityl-daunomycin conjugates (122) 107
Al-Shamkhani, A., see Morgan, S.M. (122) 121
Anwar, H., see Owusu-Ababio, G. (122) 9
Armstrong, N.A., see Moreton, R.C. (122) 79
Aspden, T.J., Adler, J., Davis, S.S., Skaugrud, Ø. and Illum, L.
Chitosan as a nasal delivery system: Evaluation of the effect of chitosan on mucociliary clearance rate in the frog palate model (122) 69

Callant, D., see Morgan, S.M. (122) 121
Casabó, V.G., see Martín-Algarra, R.V. (122) 1

Davidson, R., see Ek, R. (122) 49
Davis, S.S., see Aspden, T.J. (122) 69
Dondeti, P., Zia, H. and Needham, T.E.
In vivo evaluation of spray formulations of human insulin for nasal delivery (122) 91
Duncan, R., see Al-Shamkhani, A. (122) 107
Duncan, R., see Morgan, S.M. (122) 121

Ek, R., Lennholm, H., Davidson, R., Nyström, C. and Ragnarsson, G.
Pore swelling in beads made of cellulose fibres and fibre fragments (122) 49

Illum, L., see Aspden, T.J. (122) 69

Kanjia, J., see Pandit, N.K. (122) 27

Laughlin, S.M., see Rocco, W.L. (122) 17
Lennholm, H., see Ek, R. (122) 49

Martín-Algarra, R.V., Pascual-Costa, R.M., Merino, M. and Casabó, V.G.
Effects of polysorbate 80 on amiodarone intestinal absorption in the rat (122) 1
Merino, M., see Martín-Algarra, R.V. (122) 1
Moreton, R.C. and Armstrong, N.A.
Design and use of an apparatus for measuring diffusion through glycerogelatin films (122) 79
Morgan, S.M., Al-Shamkhani, A., Callant, D., Schacht, E., Woodley, J.F. and Duncan, R.
Alginates as drug carriers: covalent attachment of alginates to therapeutic agents containing primary amine groups (122) 121
Morphet, C., see Rocco, W.L. (122) 17
Mosharraf, M. and Nyström, C.
Solubility characterization of practically insoluble drugs using the Coulter counter principle (122) 57
Mosharraf, M. and Nyström, C.
The effect of particle size and shape on the surface specific dissolution rate of microsized practically insoluble drugs (122) 35

Needham, T.E., see Dondeti, P. (122) 91
Nyström, C., see Ek, R. (122) 49
Nyström, C., see Mosharraf, M. (122) 35
Nyström, C., see Mosharraf, M. (122) 57

Owusu-Ababio, G., Rogers, J.A. and Anwar, H.
Method of evaluation of sustained release microsphere formulations using the open chemostat system (122) 9

Pandit, N.K., Kanjia, J., Patel, K. and Pontikes, D.G.
Phase behavior of aqueous solutions containing nonionic surfactant-polyethylene glycol mixtures (122) 27
Pascual-Costa, R.M., see Martín-Algarra, R.V. (122) 1
Patel, K., see Pandit, N.K. (122) 27
Pontikes, D.G., see Pandit, N.K. (122) 27

Ragnarsson, G., see Ek, R. (122) 49
Rocco, W.L., Morphet, C. and Laughlin, S.M.
Solid-state characterization of zanoterone (122) 17
Rogers, J.A., see Owusu-Ababio, G. (122) 9

Schacht, E., see Morgan, S.M. (122) 121
Skaugrud, Ø., see Aspden, T.J. (122) 69

Woodley, J.F., see Morgan, S.M. (122) 121

Zia, H., see Dondeti, P. (122) 91